



# Attribution of the role of climate change in the forest fires in Sweden 2018

Folmer Krikken<sup>1</sup>, Flavio Lehner<sup>2</sup>, Karsten Haustein<sup>3</sup>, Igor Drobyshev<sup>4,5,6</sup>, Geert Jan van Oldenborgh<sup>1</sup>

<sup>1</sup>Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands,

<sup>2</sup>National Center for Atmospheric Research, Boulder, USA

<sup>3</sup>Environmental Change Institute, University of Oxford, Oxford, United Kingdom

<sup>4</sup>Centre for Forest Research, Montreal, Canada

<sup>5</sup>NSERC-UQAT-UQAM Industrial Chair in Sustainable Forest Management Université du Québec, Canada

<sup>6</sup>Swedish University of Agricultural Sciences, Southern Swedish Forest Research Centre, Alnarp, Sweden

Correspondence to: Folmer Krikken ([folmer.krikken@knmi.nl](mailto:folmer.krikken@knmi.nl))

**Abstract.** In this study we analyse the role of climate change in the forest fires that raged through large parts of Sweden in the summer of 2018 from a meteorological perspective. This is done by studying the Canadian Fire Weather Index (FWI) based on sub-daily data, both in reanalysis datasets (ERA-Interim, ERA5, JMA55 and MERRA2) and three large ensemble climate models (EC-Earth, W@H and CESM) simulations. The FWI based on reanalysis correlates well with observed area burned in summer ( $r=0.6$  to  $0.8$ ). We find that the maximum forest fire risk in July 2018 had return times of  $\sim 24$  years for Southern and Northern Sweden. Further, we find a negative trend of the FWI for Southern Sweden over the 1979 to 2017 time period, yielding a decreasing risk of such an event solely based on reanalysis data. However, given the short observational record, large uncertainty between the reanalysis products and large natural variability of the FWI we cannot draw robust conclusions from reanalysis data.

The 3 large-ensembles with climate models on the other hand point to a roughly 1.1 times increased risk for such events in the current climate relative to pre-industrial climate. For a future climate (2C warming) we find a roughly 2 times increased risk for such events relative to pre-industrial climate. The increased fire weather risk is mainly attributed to the increase in temperature. The other main factor, precipitation during summer months, is projected to increase for Northern Sweden, and decrease for Southern Sweden. We however do not find a clear change of prolonged dry periods in summer months that could explain the increased fire weather risk.

In summary, we find a small but positive role of global warming up to now in the 2018 forest fires in Sweden, but a more robust increase in the risk for such events in the future.



## 1 Introduction

The summer of 2018 in Sweden was characterized by numerous large forest fires spread over large parts of the country. 35 Though forest fires are common in Sweden, the number of fires and total area burned in 2018 were much higher than observed over recent years (2008-2017; Figure 1). Spring and summer weather conditions in 2018 were anomalously dry and warm. This was caused by very persistent atmospheric blocking, especially in May and July. In July the high surface pressure (Figure 2a) caused high temperature (Figure 2b) and anomalously little precipitation (Figure 2c) over northwestern Europe. The high temperature and lack of precipitation resulted in high forest fire risk over the whole of Scandinavia. 40 Especially in Sweden, this gave rise to numerous forest fires with a total burned area of more than 20.000 ha.

An often-raised question during and after such extreme events concerns the possible influence of climate change, i.e., has climate change made such an event more or less likely? Hence, climate attribution studies of extreme weather events is a rapidly increasing field of research, with analysis on e.g. extreme precipitation events (Oldenborgh et al., 2017), heat waves ( 45 Sippel et al., 2016), droughts (Hauser et al., 2017) and storms (Vautard et al., 2019), where in many cases there was indeed evidence of increased risk of extreme weather due to climate change (Schiermeier, 2018). For forest fires, the first attribution of climate change on forest fires in Canada was already found by Gillett et al. (2004) based on the CGCM2 model. Abatzoglou and Williams (2016) found that for the western United States human-caused climate change more than halved humidity of forest fuels since the 1970s and doubled the cumulative area of forest fires since 1984. A recent study by 50 Kirchmeier-Young et al., (2018) found a strong influence of climate change on the 2017 British Columbia wildfires, with such events being 2-4 times more likely with climate change in the CanESM2 model. Abatzoglou et al. (2018) found that increases in extreme fire weather days due to anthropogenic climate change are evident on 22% of burnable land area globally.

55 For Sweden specifically, Yang et al. (2015) found that in a future climate there is an increased risk of forest fires in Southern Sweden but a decreased risk in Northern Sweden using the downscaled and bias corrected ECHAM5 climate model. Also, for the neighbouring country of Finland, climate model projections point to an increased risk of forest fires (Lehtonen et al., 2016).

60 Here we analyse the connection between the 2018 extreme forest fire season and climate change using large ensembles of multiple climate models. As characteristics of regional precipitation and droughts can be highly model dependent (Hauser et al., 2017), it is crucial to use multiple climate models for this analysis. To the best of our knowledge, this is the first such multi-model framework applied to an attribution study of an extreme forest fire event.



65 Note that we only analyse the meteorological aspect of this event, and not other aspects relevant for such extreme events  
such as the sources of ignition and the influence of fire mitigation strategies. Hence, in our analysis we take forest fire risk to  
be equivalent to fire weather risk. We do investigate to what extent the fire risk corresponds to actual area burnt.

## 2 Data and Methods

### 70 2.1 Fire weather risk

The metric used to quantify forest fire risk is the Canadian Fire Weather Index (FWI; Van Wagner, 1987). This is a weather-  
based system that models soil moisture at 3 different depths, and, based on the upper soil moisture content and wind speed,  
creates an estimate for the initial spread rate of fire. It is based on four meteorological variables, namely local noon  
temperature (T2M), relative humidity (RH), surface wind speed and 24 hour cumulative precipitation. Though this metric  
75 was developed and tuned for the Canadian boreal region, it also performs well over Sweden (Gardelin, 1997; Yang et al.,  
2015).

### 2.2 Statistical methods

In event attribution studies, the first step is to define the event in such a way that is best reflects the impact of the event. In  
80 section 3 we will discuss how we define this event in more detail. The second step, in order to assess the rarity of this event,  
is to fit a Generalized Extreme Value Distribution (GEV) function on a sample of block (yearly) maxima extracted from a  
FWI time series. The GEV function is described by three parameters: the position parameter  $\mu$ , the scale parameter  $\sigma$  and the  
shape parameter  $\xi$ .

85 In order to assess the risks of certain events during previous climate based on the trend in observations and reanalysis  
products, we fit the observed FWI to a GEV that depends on the smoothed (4th order polynomial) global mean surface  
temperature (GMST). Here, GMST is taken from the National Aeronautics and Space Administration (NASA) Goddard  
Institute for Space Science (GISS) surface temperature analysis (GISTEMP), (Hansen et al., 2010). This results in a  
distribution that varies continuously with GMST. This distribution can be evaluated for a GMST in the past (e.g., 1950 or  
90 1900) and for the current GMST. A 1000-member non-parametric bootstrap procedure is used to estimate confidence  
intervals for the fit.

For the FWI we choose the dependence of the FWI the same way as for precipitation (described fully in Wiel et al., 2017):  
the position and scale parameters ( $\mu, \sigma$ ) have the same dependence on GMST so that their ratio (also called the dispersion



95 parameter) is constant. The dependence is exponential for precipitation,  $\mu(T) = \mu_0 \exp(\alpha T / \mu_0)$ ,  $\sigma(T) = \sigma_0 \exp(\alpha T / \mu_0)$ , we use the same here.

We use the risk ratio to quantify the impact of climate change on the FWI. This ratio, calculated as the risk of an event occurring during the current or future climate divided by the risk of an event occurring during pre-industrial climate conditions, indicates how much more or less likely a certain event will be relative to pre-industrial climate. Thus, a risk ratio of 2 means an event will be 2 times more likely relative to pre-industrial climate.

### 2.3 Reanalysis

We use multiple reanalysis dataset as an estimate of the observed state, namely ERA-Interim (ERA-I, Dee et al., 2011), ERA5 (Copernicus Climate Change Service, 2017), the Japanese 55-year Reanalysis (JRA-55, Kobayashi et al., 2015) and the Modern-Era retrospective analysis for Research and Application, Version 2 (MERRA2, Global Modeling and Assimilation Office, 2015). The advantage of using reanalysis datasets compared to in-situ observations is that it provides an observationally constrained continuous gridded dataset, enabling direct comparison to climate model output. The orography of Sweden is such that the relatively low-resolution models used to generate the reanalysis can represent the weather well in this area. We use multiple reanalysis products in order to sample the uncertainty in these products. All products provide a continuous dataset from 1979/1980 to current, with the exception of JRA-55 which spans the period from 1955 to current.

### 2.4 Models

We use climate model simulations from three different coupled climate models with large ensembles: EC-Earth v2.3 (Hazeleger et al., 2010; 2011), the Community Earth System Model version 1 (CESM1, (Kay et al., 2014) and Weather@Home (W@H, Guillod et al., 2017; Massey et al., 2015) (Table 1). The EC-Earth and CESM are large ensembles of transient climate simulations with historical forcing prior to 2006 and the RCP85 forcing (Riahi et al., 2011) from 2006 onwards. The W@H climate simulations are two different simulation, one with the actual observed forcing to represent current climate, and one with natural forcing only (i.e. no anthropogenic forcing) to represent pre-industrial climate. From EC-EARTH and CESM we select three periods that (1) describe the unperturbed climate (i.e., pre-industrial climate), (2) the current climate and (3) the 2°C warming threshold (future climate). **The pre-industrial, current and future climate states are hereafter referred to as respectively PI, 1C and 2C.** Note that ‘current climate’, or ‘1C’, is chosen in a way to best compare to reanalysis (which covers the years 1979-2018) as described below.

125 First, we select the time periods from EC-Earth and CESM1 that represent the same incremental global warming from ‘PI’ to ‘1C’ as in observations: in GISTEMP, GMST increases by 0.67°C between 1900-1950 and 1979-2018. Finding the same



130 warming increment in EC-EARTH and CESM results in the time periods listed in Table 1 for the two models. For the ‘2C’ climate we select a 30 year window with a 2°C warming relative to PI (Table 1). For the W@H simulations the GMST increase between the ‘natural forcing’ simulations and the ‘actual forcing’ simulations is 0.65<sup>0</sup>C, which is very close to the observed warming.

135 A second bias correction step is performed on the basis of return times of the specific event and can be seen as a local bias correction in contrast to the first step, which aimed at aligning simulations and observations with regard to the level of global warming. We first calculate the return time of the event from observations or reanalysis (Figure 3). In the ‘1C’ model simulations we then select the FWI that corresponds to that specific return time. This FWI value is used to estimate return times in the other simulated climate states (PI and 2C). The advantage of this approach is that it preserves the spatiotemporal consistency of the simulated fields, the relation among the meteorological variables, and it makes no assumptions on non-stationarity in bias correction, which are typical issues in (multi-variate) bias correction methods (Ho et al. 2011; Ehret et al. 2012). Note that this bias correction is only a viable method if the risk ratio is not too sensitive to the event return time, 140 which is the case here (not shown).

We calculate the FWI on the original grid of the models. Since Southern Sweden has a different fire climate than Northern Sweden (Drobyshev et al., 2012), we calculate spatial averages for northern (Norrland), middle (Svealand) and southern (Göteborg) Sweden (Figure 4a). Since high fire weather risk events are mainly associated with large high pressure systems, it is important to validate the persistence of high pressure systems in the climate models by comparing it to reanalysis data. 145 Following the method of Pfleiderer and Coumou (2018), which represents persistence as the number of consecutive warm days, we find that the models are in good agreement with reanalysis in respect to persistence of high pressure systems (not shown).

150 For EC-Earth we compute the FWI based on local noon data (12 UTC), but for the CESM-LE and W@H sub-daily data is not available. Hence, for these models we compute the FWI based on daily average wind speed and humidity, daily maximum temperature and daily cumulative precipitation. While this approach is common (Abatzoglou et al., 2019), results can differ between both methods especially for fire danger extremes (Herrera et al., 2013). In order to assess whether this has an effect on our analysis we tested the influence of using local noon data, or daily average combined with maximum 155 temperature for EC-Earth. Though the values of the FWI do differ, there are no significant differences for both methods on the calculated risk ratios. Hence, we assume that using daily maximum temperature and daily average values for the other variables for the calculation of the FWI in CESM-LENS and W@H does not affect the calculated risk ratios significantly.



### 3 Event definition

We first investigate whether the FWI is a good proxy for actual fires in Sweden. For the event definition we use ERA-  
160 Interim as the observational estimate. The FWI is a physical approximation of climatological fire risk and it has been found  
to be a robust proxy for actual fires (Wotton, 2009). However, there can be a strong seasonal dependence on the correlation  
between the FWI and actual fires (Lehtonen et al., 2016). We test this for Sweden by studying the correlation between the  
FWI and observed area burned (MSB, 2016) for the period 1996 to 2012 (Figure 4b). Note that, here we leave out Svealand  
because there are insufficient fires to compute the relevant statistics. The results show that there is indeed a strong seasonal  
165 dependence on the relation between monthly averaged FWI and area burned, with generally high correlations from July  
onwards, but lower correlations for April to June for Norrland and May for Götaland. These findings correspond to the  
findings from Lehtonen et al. (2016), who relate the low correlation in spring to the influence of more human-caused fires,  
whilst in summer natural ignitions is a more important ignition source thus yielding a stronger relation with weather  
variables.

170

Next, we analyse the FWI for all three regions, using ERA-Interim (Figure 5) and the observed area burned (Figure 1). There  
were two distinct periods of high FWI (above the 95% quantile), namely in late May to early June and in July. Interestingly,  
the values in May were even higher than those observed in July, although the actual area burned was much higher in July.  
This indicates a possible pre-conditioning (drying out the soil) of summer FWI by the occurrence of a dry spring. Note that  
175 the pre-conditioning by the dry spring is still included in this event definition because the FWI calculation includes an  
estimate of the moisture content in the deeper soil layer. This moisture content, estimated by the Drought Code within the  
FWI calculation, includes memory of ~52 days (Van Wagner, 1987).

Based on the findings from Figure 1, 4 and 5, we define our event as the maximum 7-day running mean FWI in the months  
180 of July and August, disregarding the FWI peak in May due to much lower correlations with area burned, probably due to  
lower ignition rates. The 7-day running mean is applied as fires are more likely to happen during prolonged period of high  
fire weather risk, whilst still holding enough independent samples per year for a robust GEV fit. Though June also shows  
relative high correlation with observed area burned, the strong fires were mainly in the summer months.

## 4. Results

### 185 4.1 Reanalysis

As previously stated, July 2018 was characterized by a large persistent high pressure area over Northern Europe (Figure 2),  
yielding high temperatures, little precipitation and moderate winds. The meteorological conditions for fire weather were thus  
quite extreme. This is quantified by the high return times of such conditions for July 2018 (Figure 6). These values are based



on a GEV-fit, based on the maximum value of FWI in July and August for every year, with a 7-day running mean applied.  
190 This fit assumes that the climate does not change over time.

It is striking to see that, although all reanalysis products are constrained by observations, there are still quite large differences  
in the FWI value for the 2018 event and the associated return times. For Norrland, we find large significant differences  
between JMA-55, with a return time of ~5 yr, and ERA-I, ERA5 and MERRA-2 with return times of ~30 yr. Also for  
195 Svealand there are rather large differences in return times, with ~10 years for JMA-55 and ~20 yr for ERA-I and ERA5 and  
~50 yr for MERRA-2. In Götaland we also find differences between the products, but now JMA-55 closely matches ERA5  
with return times of ~8-10 yr, and ERA-I has a higher return times of ~20 yr and MERRA-2 even higher at ~60 yr. Note that  
the uncertainties on these return times (denoted by the horizontal bars) are large but almost completely correlated across  
datasets as they derive from the same natural variability (except JMA-55 that includes more data). An analysis on the  
200 meteorological variables used in the FWI reveals that it is mainly precipitation that causes the differences in FWI and return  
times across products. For MERRA2, it is also related to a generally lower temperature (not shown).

These results stress the importance of using multiple reanalysis products in order to get a better estimate of the observed  
event and its associated uncertainty. For bias correction of the climate model data, we use the average of all four return times  
205 from the different reanalysis products.

In order to analyse whether such an event has become more or less likely relative to a climate without anthropogenic  
emissions ("pre-industrial", PI), we fit the yearly maximum FWI to a GEV that scales with the smoothed GMST (as  
described in the methods section). We can then evaluate the probability of such an event conditional on different climate  
210 states as defined by GMST. Figure 7a shows the risk ratios for the reanalysis products. Note that, as stated before, the risk  
ratio tells us how much more or less likely such an event has become today ('1C' climate) relative to PI climate. The  
reanalysis data shows a slightly decreased risk of high FWI events for all three regions for the 1C climate. This is due to a  
negative trend of July and August FWI over recent decades for these regions. However, the very large uncertainties easily  
encompass one (no change), indicating that this is most likely a spurious trend caused by natural variability. This trend is  
215 largely absent in JMA-55 which has a 25 year longer time span than the other reanalysis datasets. Hence, we cannot draw  
any robust conclusions from the trends of the reanalysis dataset alone.

## 4.2 Models

With large ensemble climate model data we can circumvent the problems of undersampled natural variability, allowing us to  
220 get more robust estimates of whether the likelihood of such an event changes with time. Figure 7b shows the risk ratios of



the climate models for present climate (1C) relative to PI climate and future climate (2C) relative to PI climate. First we will focus on the comparison of 1C to PI.

The model W@H shows a small, but significant increased risk of approximately a factor 1.5 for such events for all three regions. EC-Earth shows no clear change in risk for such an event, with risk ratios close to 1, whereas CESM does show a small increase in risk, though not significant. **On average, we find a small (not significant) increase in risk for all three regions.** In the 2C climate, the risk ratios increase more strongly relative to PI climate. CESM shows significant increased risk ratios of ~3, with the largest increase in risk in Norrland. EC-Earth also shows an increased risk, though not as large as CESM. On average for all three regions we find a risk ratio of ~2

## 230 5 Discussion

In general, **we find a factor 1.2 increased risk for such events for current climate relative to PI climate,** and a significant increase in risk of factor 2 to 3 for a 2°C warmer climate relative to PI climate. To better understand why there is an increased risk of such events, we investigate the individual meteorological variables at the time of the maximum July and August FWI in models (Figure 8).

235

All models show a clear trend towards higher temperatures, which is unsurprising as present day and future climate are chosen as ~1°C and 2°C warmer climates. The increase in temperature between 1C and 2C is generally much larger than 1°C, especially in CESM-LE under 2C, because land heats up faster than the global mean. This can partly explain the relative strong increase in fire risk in CESM-LE for future climate. In EC-Earth the relative humidity seems to reflect the changes in precipitation where it increases from PI to 1C and then decreases slightly in the 2C climate. In CESM-LE, we find no clear change in 1C, but a decrease of RH in 2C. W@H also show a small decrease in RH relative to PI climate. For the wind speed, the differences between the climate states are very small and do not affect changes in the FWI appreciably. We further subset the model values by focusing on FWI events larger than the 2018 observed event (circles in Fig. 8) to investigate whether certain variables are predominantly affecting extreme FWI. We find no relationship between wind and extreme FWI values, indicating wind is not an important explanatory variable for extreme FWI events over Sweden.

245

For precipitation, we generally find an increase from PI to 1C for all regions and climate models, with the exception of W@H with a small decrease in precipitation. For 2C, however, there are strong differences between the regions, where in Norrland precipitation further increases whilst for Svealand and Götaland it decreases towards PI values (EC-Earth) or stays constant with present climate values (CESM-LE). Note however, that the precipitation values associated with high FWI values (circles in Figure 8) do not show this upward trend for Norrland. Hence, changes in mean precipitation do not necessarily reflect the changes in prolonged dry periods. An analysis of the trends in the lower (dry periods) and middle quantiles of 30-day precipitation in summer shows clear changes in the median but no clear changes in the lower quantiles

250





(not shown). This is also demonstrated by Pendergrass et al. (2017), who find that precipitation variability generally  
255 increases in a warmer climate.

The changes in FWI between the different climates relate mostly to changes in precipitation and temperature, as RH follows  
the changes of these variables. The higher temperatures for present and future climate relative to PI climate yield an increase  
of FWI. However, the increase of precipitation can counteract this increase. The fact that FWI increases even in models with  
260 mean increases in precipitation shows that temperature increases dominate future increases in FWI.

One potential reason for the non-linear model behaviour with regard to the different warming thresholds is the non-linear  
evolution of the radiative forcing. Global dimming associated with the release of cooling anthropogenic aerosol particles  
(Wild 2009) has effectively offset the greenhouse gas induced warming during 1950-1975. In contrast, global brightening  
265 has accelerated the warming thereafter, particularly over North America and Europe where sulphate aerosols emissions were  
curbed substantially. As a result, aerosols may have suppressed rainfall during the dimming phase, but invigorated  
precipitation during the subsequent warming until now, exceeding the direct temperature effect. The smaller risk reduction in  
the longer JMA-55 time series would support this hypothesis.

270 Another dynamic factor that is projected to come into play over the next decades is the development of a heat low over the  
Mediterranean area. This would increase the possibility of easterly wind over northern Europe and hence dry weather,  
offsetting the trend towards wetter summer weather up to now (Haarsma et al., 2009).

Our results mostly agree with previous research. The work of Flannigan et al. (2012) points to increased risk of forest fires  
275 over the whole of Sweden for multiple climate projections. Findings from Yang et al. (2015) point to an increased risk of  
forest fires in the southern part of Sweden, but not for the northern part where they point to increased precipitation which  
reduces fire risk. This difference can be caused by undersampling of extreme events, since Yang et al use a single 30 year  
time slice of future (2071-2100) climate. It must also be noted that local future precipitation trends are highly uncertain  
(Lehtonen et al., 2016), implying that using only one climate model for future projection (as in Yang et al., 2016). Even the  
280 three climate models in this study will likely underestimate the model uncertainty in the precipitation trends. Hence, future  
work should focus on using more large ensemble climate models in order to better sample the uncertainty in the future  
climate projections.

Note we assume that FWI remains a skillful predictor of area burned (Figure 4), even in a future climate. This assumption is  
285 however highly uncertain due to factors not accounted for in the analysis here, such as possible changes in forest  
management (Moreira and Pe'er, 2018; Hudson, 2018), a possible increase / decrease of human-caused forest fires (Balch et  
al., 2017) and feedback mechanisms between forest fires and ecology (Balch et al., 2008).



## 6. Conclusions

290

In our analysis of the forest fires in Sweden of 2018 we have looked at the risk of fire weather solely on the basis of the Canadian FWI, with the novel approach of using multiple reanalysis datasets and multiple large ensembles with climate models. Using the FWI we have only attributed meteorological aspects of this event, but acknowledge that there are additional aspects important for determining forest fire risk not considered here.

295

We find that the maximum forest fire risk in July 2018 had return times of ~24 years in Götaland, ~23 years in Svealand and ~24 years in Norrland, with large uncertainty in the reanalysis datasets (90% uncertainty estimate  $\pm$  ~10 years). Due to the relative short observational record, large uncertainty in the reanalysis datasets and large natural variability of the FWI we cannot infer a robust trend from the reanalysis data alone.

300

The climate models point to a small increase in risk for such an event at present day compared to pre-industrial conditions for all three regions of about a factor 1.1 (0.9 to 1.4). In a future climate (a 2<sup>o</sup>C warmer climate relative to pre-industrial) the risk for such events to occur may increase more robustly by a factor of ~2 (1.5 to 3) relative to pre-industrial climate according to our model analysis.

305

The increased fire risk is mostly driven by increased temperature. Though we do find clear changes in precipitation for the warmer climates, we do not see a clear change in prolonged dry periods during summer, which have historically and will likely continue to drive high fire risk events. Our results show the importance of using multiple large ensembles with climate models for attribution studies in order to adequately sample the natural variability and model uncertainties in climate

310

projections.



### Code availability

The Python code used to compute the FWI and for the analysis can be obtained by contacting Folmer Krikken  
315 ([folmer.krikken@knmi.nl](mailto:folmer.krikken@knmi.nl))

### Data availability

EC-Earth data can be obtained by contacting Folmer Krikken ([folmer.krikken@knmi.nl](mailto:folmer.krikken@knmi.nl)), CESM data can be obtained by  
contacting Flavio Lehner ([flehner@ucar.edu](mailto:flehner@ucar.edu)) and W@H data can be obtained by contacting Karsten Haustein  
320 ([karsten.haustein@ouce.ox.ac.uk](mailto:karsten.haustein@ouce.ox.ac.uk)).

### Author contribution

G.J.O. and F.K. conceived and planned the analysis. F.K. provided the EC-Earth data. F.L. provided the CESM data. K.H.  
provided the W@H data. F.K. performed the main analysis and wrote the main manuscript with support from all authors at  
325 all stages.

### Competing interests

The authors declare that they have no conflict of interests

### 330 Acknowledgements

F.K. and I.D. contribution was supported by the Belmont Forum Project PRERREAL (grant # 292-2015-11-30-13-43-09 to  
I.D.). I.D. contribution was further supported by the Canadian National Research Council Canada through Discovery Grant  
(grant # DDG-2015-00026 to I.D.), Swedish Research Council FORMAS (grant # 239-2014-1866 to I.D.), The Swedish  
Institute funded networks CLIMECO and BalticFire (#10066-2017-13 and #24474/2018 to I.D. The study was conducted  
335 within the framework of the NordicProxy network, which is supported by the Nordic Forest Research (SNS), and consortium  
GDRI Cold Forests. G.J.O. was supported by the ERA4CS project SERV\_FORFIRE. F.L. is supported by NSF AGS-  
0856145, Amendment 87, by the Bureau of Reclamation under Cooperative Agreement R16AC00039, and the Regional and  
Global Model Analysis (RGMA) component of the Earth and Environmental System Modeling Program of the U.S.  
Department of Energy's Office of Biological & Environmental Research (BER) Cooperative Agreement DE-FC02-  
340 97ER62402. We thank Guilherme S.J. Pinto for providing the Swedish forest fire data from the MSB.



## References

- 345 Abatzoglou, J., Williams, A. and Barbero, R.: Global Emergence of Anthropogenic Climate Change in Fire Weather Indices, *Geophys. Res. Lett.*, doi:10.1029/2018GL080959, 2018.
- Abatzoglou, J. T. and Williams, A. P.: Impact of anthropogenic climate change on wildfire across western US forests, *Proc. Natl. Acad. Sci.*, 113(42), 11770–11775, doi:10.1073/pnas.1607171113, 2016.
- Balch, J. K., Nepstad, D. C., Brando, P. M., Curran, L. M., Portela, O., Carvalho, O. D. and Lefebvre, P.: Negative fire  
350 feedback in a transitional forest of southeastern Amazonia, *Glob. Change Biol.*, 14(10), 2276–2287, doi:10.1111/j.1365-2486.2008.01655.x, 2008.
- Balch, J. K., Bradley, B. A., Abatzoglou, J. T., Nagy, R. C., Fusco, E. J. and Mahood, A. L.: Human-started wildfires expand the fire niche across the United States, *Proc. Natl. Acad. Sci.*, 114(11), 2946–2951, doi:10.1073/pnas.1617394114, 2017.
- 355 Copernicus Climate Change Service (C3S) (2017): ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store (CDS), *Date accessed: 15-03-2019*.  
<https://cds.climate.copernicus.eu/cdsapp#!/home>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R.,  
360 Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828, 2011.
- Drobyshev, I., Niklasson, M. and Linderholm, H. W.: Forest fire activity in Sweden: Climatic controls and geographical  
365 patterns in 20th century, *Agric. For. Meteorol.*, 154–155, 174–186, doi:10.1016/j.agrformet.2011.11.002, 2012.
- Ehret, U., Zehe, E., Wulfmeyer, V., Warrach-Sagi, K. and Liebert, J.: HESS Opinions “Should we apply bias correction to global and regional climate model data?,” *Hydrol. Earth Syst. Sci.*, 16(9), 3391–3404, doi:https://doi.org/10.5194/hess-16-3391-2012, 2012.
- Gardelin, M.: Brandriskprognoser med hjälp av en kanadensisk skogsbrandsmodell, Räddningsverket Report, Myndigheten  
370 för samhällsskydd och Beredskap (MSB), Sweden, 1997.
- Gillett, N. P., Weaver, A. J., Zwiers, F. W. and Flannigan, M. D.: Detecting the effect of climate change on Canadian forest fires, *Geophys. Res. Lett.*, 31(18), doi:10.1029/2004GL020876, 2004.
- Global Modeling and Assimilation Office (GMAO) (2015), MERRA-2 inst3\_3d\_asm\_Nv: 3d,3-Hourly,Instantaneous,Model-Level,Assimilation,Assimilated Meteorological Fields V5.12.4, Greenbelt, MD, USA,



- 375 Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: 15-03-2019,  
10.5067/WWQSQ8IVFW8
- Guillod, B. P., Jones, R. G., Bowery, A., Haustein, K., Massey, N. R., Mitchell, D. M., Otto, F. E. L., Sparrow, S. N., Uhe,  
P., Wallom, D. C. H., Wilson, S. and Allen, M. R.: weather@home 2: validation of an improved global–regional  
climate modelling system, *Geosci. Model Dev.*, 10(5), 1849–1872, doi:https://doi.org/10.5194/gmd-10-1849-2017,  
380 2017.
- Haarsma, R. J., Selten, F., Hurk, B. vd, Hazeleger, W. and Wang, X.: Drier Mediterranean soils due to greenhouse warming  
bring easterly winds over summertime central Europe, *Geophys. Res. Lett.*, 36(4), doi:10.1029/2008GL036617,  
2009.
- Hansen, J., Ruedy, R., Sato, M. and Lo, K.: Global Surface Temperature Change, *Rev. Geophys.*, 48(4),  
385 doi:10.1029/2010RG000345, 2010.
- Hauser, M., Gudmundsson, L., Orth, R., Jézéquel, A., Haustein, K., Vautard, R., Oldenborgh, G. J. van, Wilcox, L. and  
Seneviratne, S. I.: Methods and Model Dependency of Extreme Event Attribution: The 2015 European Drought,  
*Earths Future*, 5(10), 1034–1043, doi:10.1002/2017EF000612, 2017.
- Hazeleger, W., Severijns, C., Semmler, T., Ștefănescu, S., Yang, S., Wang, X., Wyser, K., Dutra, E., Baldasano, J. M.,  
390 Bintanja, R., Bougeault, P., Caballero, R., Ekman, A. M. L., Christensen, J. H., van den Hurk, B., Jimenez, P.,  
Jones, C., Kållberg, P., Koenigk, T., McGrath, R., Miranda, P., Van Noije, T., Palmer, T., Parodi, J. A., Schmith,  
T., Selten, F., Storelvmo, T., Sterl, A., Tapamo, H., Vancoppenolle, M., Viterbo, P. and Willén, U.: EC-Earth: A  
Seamless Earth-System Prediction Approach in Action, *Bull. Am. Meteorol. Soc.*, 91(10), 1357–1363,  
doi:10.1175/2010BAMS2877.1, 2010.
- 395 Hazeleger, W., Wang, X., Severijns, C., Ștefănescu, S., Bintanja, R., Sterl, A., Wyser, K., Semmler, T., Yang, S., Hurk, B.  
van den, Noije, T. van, Linden, E. van der and Wiel, K. van der: EC-Earth V2.2: description and validation of a new  
seamless earth system prediction model, *Clim. Dyn.*, 39(11), 2611–2629, doi:10.1007/s00382-011-1228-5, 2011.
- Herrera, S., Bedia, J., Guti rrez, J. M., Fern ndez, J. and Moreno, J. M.: On the projection of future fire danger conditions  
with various instantaneous/mean-daily data sources, *Clim. Change*, 118(3), 827–840, doi:10.1007/s10584-012-  
400 0667-2, 2013.
- Ho, C. K., Stephenson, D. B., Collins, M., Ferro, C. A. T. and Brown, S. J.: Calibration Strategies: A Source of Additional  
Uncertainty in Climate Change Projections, *Bull. Am. Meteorol. Soc.*, 93(1), 21–26,  
doi:10.1175/2011BAMS3110.1, 2011.
- Hudson, B.: Fighting Fire with Fire? Adjusting Regulatory Regimes and Forest Product Markets to Mitigate Southern United  
405 States Wildfire Risk, SSRN Scholarly Paper, Social Science Research Network, Rochester, NY. [online] Available  
from: <https://papers.ssrn.com/abstract=3179338> (Accessed 30 January 2019), 2018.
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., Edwards,  
J., Holland, M., Kushner, P., Lamarque, J.-F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R.,



- Oleson, K., Polvani, L. and Vertenstein, M.: The Community Earth System Model (CESM) Large Ensemble Project: A Community Resource for Studying Climate Change in the Presence of Internal Climate Variability, *Bull. Am. Meteorol. Soc.*, 96(8), 1333–1349, doi:10.1175/BAMS-D-13-00255.1, 2014.
- Kirchmeier-Young, M. C., Gillett, N. P., Zwiers, F. W., Cannon, A. J. and Anslow, F. S.: Attribution of the Influence of Human-Induced Climate Change on an Extreme Fire Season, *Earths Future*, 0(0), doi:10.1029/2018EF001050, 2018.
- 415 Kobayashi, S., Ota, Y., Harada, Y., Ebita, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K. and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *J. Meteorol. Soc. Jpn. Ser II*, 93(1), 5–48, doi:10.2151/jmsj.2015-001, 2015.
- Lehtonen, I., Venäläinen, A., Kämäräinen, M., Peltola, H. and Gregow, H.: Risk of large-scale fires in boreal forests of Finland under changing climate, *Nat. Hazards Earth Syst. Sci.*, 16(1), 239–253, doi:https://doi.org/10.5194/nhess-16-239-2016, 2016.
- 420 Massey, N., Jones, R., Otto, F. E. L., Aina, T., Wilson, S., Murphy, J. M., Hassell, D., Yamazaki, Y. H. and Allen, M. R.: weather@home—development and validation of a very large ensemble modelling system for probabilistic event attribution, *Q. J. R. Meteorol. Soc.*, 141(690), 1528–1545, doi:10.1002/qj.2455, 2015.
- Moreira, F. and Pe'er, G.: Agricultural policy can reduce wildfires, *Science*, 359(6379), 1001–1001, doi:10.1126/science.aat1359, 2018.
- 425 MSB, 2017. Incident reports from municipal fire brigades. Swedish Civ. Conting. Agency (Myndigheten för samhällsskydd och Beredsk in Swedish). <https://www.msb.se/>
- Oldenborgh, G. J. van, Wiel, K. van der, Sebastian, A., Singh, R., Arrighi, J., Otto, F., Haustein, K., Li, S., Vecchi, G. and Cullen, H.: Attribution of extreme rainfall from Hurricane Harvey, August 2017, *Environ. Res. Lett.*, 12(12), 124009, doi:10.1088/1748-9326/aa9ef2, 2017.
- 430 Pendergrass, A. G., Knutti, R., Lehner, F., Deser, C. and Sanderson, B. M.: Precipitation variability increases in a warmer climate, *Sci. Rep.*, 7(1), 17966, doi:10.1038/s41598-017-17966-y, 2017.
- Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N. and Rafaj, P.: RCP 8.5—A scenario of comparatively high greenhouse gas emissions, *Clim. Change*, 109(1), 33, doi:10.1007/s10584-011-0149-y, 2011.
- 435 Schiermeier, Q.: Droughts, heatwaves and floods: How to tell when climate change is to blame, *Nature*, 560, 20, doi:10.1038/d41586-018-05849-9, 2018.
- Sippel, S., Otto, F. E. L., Flach, M. and van Oldenborgh, G. J.: The Role of Anthropogenic Warming in 2015 Central European Heat Waves, *Bull. Am. Meteorol. Soc.*, 97(12), S51–S56, doi:10.1175/BAMS-D-16-0150.1, 2016.
- 440 Van Wagner, C. E.: Development and structure of the Canadian forest fire index system, *Can. For. Serv.*, 35, 1987.
- Vautard, R., Oldenborgh, G. J. van, Otto, F. E. L., Yiou, P., Vries, H. de, Meijgaard, E. van, Stepek, A., Soubeyroux, J.-M., Philip, S., Kew, S. F., Costella, C., Singh, R. and Tebaldi, C.: Human influence on European winter wind storms



- such as those of January 2018, *Earth Syst. Dyn.*, 10(2), 271–286, doi:<https://doi.org/10.5194/esd-10-271-2019>, 2019.
- 445 Wiel, K. van der, Kapnick, S. B., Oldenborgh, G. J. van, Whan, K., Philip, S., Vecchi, G. A., Singh, R. K., Arrighi, J. and Cullen, H.: Rapid attribution of the August 2016 flood-inducing extreme precipitation in south Louisiana to climate change, *Hydrol. Earth Syst. Sci.*, 21(2), 897–921, doi:<https://doi.org/10.5194/hess-21-897-2017>, 2017.
- Wild, M.: Global dimming and brightening: A review, *J. Geophys. Res. Atmospheres*, 114(D10), doi:10.1029/2008JD011470, 2009.
- 450 Wotton, B. M.: Interpreting and using outputs from the Canadian Forest Fire Danger Rating System in research applications, *Environ. Ecol. Stat.*, 16(2), 107–131, doi:10.1007/s10651-007-0084-2, 2009.
- Yang, W., Gardelin, M., Olsson, J. and Bosshard, T.: Multi-variable bias correction: application of forest fire risk in present and future climate in Sweden, *Nat. Hazards Earth Syst. Sci.*, 15(9), 2037–2057, doi:<https://doi.org/10.5194/nhess-15-2037-2015>, 2015.


455



460 **Table 1: Overview of the climate models and the years used to represent the different climate states.**


<b>Model</b>	<b>members</b>	<b>Past climate (PI)</b>	<b>Current climate (1°C)</b>	<b>Future climate (2<sup>0</sup>C)</b>	<b>Resolution</b>
<b>EC-Earth</b>	16	1900-1950 800 yrs total	1979-2019 640 yrs total	2029-2059 480 yrs	1.1°
<b>CESM1</b>	40	1920-1950 1200 yrs total	1987-2027 1600 yrs total	2028-2058 1200 yrs	1°
<b>W@H</b>	100	Natural forcing 1986-2015 3000 yrs total	Actual forcing 1986-2015 3000 yrs total	Not available	0.25°






465

Figure 1: Area burned in Sweden. Cumulative values for 2018 and climatology over 2008-2017 and its individual years (source: EFFIS).




470 **Figure 2: ERA-Interim July average anomalies of a) mean sea level pressure (MSL), b) surface temperature and c) precipitation. Anomalies are constructed relative to 1981-2010 climatology**




**Figure 3: Schematic of the bias correction method.**

475




480


Figure 4: a) Map of Sweden with the three regions used in this study and b) correlation of FWI (ERA-Interim, monthly maximum value with a 7-day running mean applied) with observed area burned for Norrländ and Götaland from 1998 to 2017. The dotted lines represent the 5-95% bootstrapped confidence intervals and the gray line indicates the significance threshold of 5%. The observed area burned is from Swedish governmental data (MSB, 2016).




485 **Figure 5: Area averaged FWI for the three regions (defined in Figure 3). The (thick) red line shows the (7-day running mean) FWI of 2018. The black lines represent the 5, 50 and 95% quantiles of the 1979-2017 climatology and the opaque gray lines the individual years, all based on ERA-Interim extended with ECMWF forecast analysis.**



490 **Figure 6: Return times of July–August maximum FWI values, for all four reanalysis datasets and the three regions. The dots represent the actual FWI maximum values and the lines the GEV model fit with a 5 to 95% uncertainty band in gray. The dashed horizontal lines represents the 2018 event, whilst the vertical line represents the associated return time with the horizontal bars giving the 5% to 95% uncertainty estimate (estimated with a non-parametric bootstrap).**



495 **Figure 7: Risk ratios for maximum July-August FWI values as high as observed in 2018 for the different regions for a) reanalysis and b) climate models. All risk ratios are relative to PI climate. Note the different scales on the x-axis between (a) and (b).**



500 **Figure 8: Meteorological values associated with the yearly maximum FWI in July and August, with a 7-day rolling average applied, for all three climate states, all three regions. Precipitation is calculated as 30 day cumulative value prior to the yearly maximum FWI. The boxplot shows the quartiles of the distribution, the whiskers the rest of the distribution and the dots are outliers. The round circles indicate all values in the distribution associated with FWI higher than the observed 2018 event.**