

The role of surface heat fluxes in tropical intraseasonal oscillations

The tropics sustain strong, coherent variations in wind and precipitation on the intraseasonal (30–60 day) timescale. In their active phases, these intraseasonal oscillations are characterized by the slow eastward movement of stronger-than-average precipitation and westerly winds. In northern summer, rainfall and wind anomalies also propagate northward on the intraseasonal timescale over India, southeast and east Asia and the adjacent oceans, pacing the active and break cycles of the monsoons and thus exerting a direct control on the livelihoods of large populations dependent on rain-fed agriculture. We argue that heat fluxes from ocean to atmosphere play a fundamental role in driving the intraseasonal oscillations. We also propose that the current generation of numerical models may enable us to test this and other hypotheses about the dynamics of intraseasonal oscillations more convincingly than has been done in the past.

ADAM H. SOBEL^{1*}, ERIC D. MALONEY²,
GILLES BELLON³ AND DARGAN M. FRIERSON⁴

¹Department of Applied Physics and Applied Mathematics and Department of Earth and Environmental Sciences Columbia University, 500 West 120th Street, Rm. 217, New York 10027, USA

²Department of Atmospheric Sciences, Colorado State University, Fort Collins, Colorado 80523, USA

³Department of Applied Physics and Applied Mathematics, Columbia University, New York 10027, USA

⁴Department of Atmospheric Sciences, University of Washington, Seattle, Washington 98195, USA

*e-mail: ahs129@columbia.edu

In addition to their direct control on local weather, the intraseasonal oscillations (ISOs) have several indirect impacts. They modulate the occurrence of tropical cyclones, not only in the Indian and western Pacific ocean basins but also in the eastern Pacific and Atlantic^{1,2}, influence weather at extratropical latitudes^{3,4} and may also play a role in initiating El Niño events⁵. Prediction of ISOs is thus of great value and has recently begun to be done with some success, using both statistical methods and numerical models^{6–8}. Recent intercomparison studies^{9,10} show that the state of the art in climate models' ISO simulations has improved significantly compared with a decade earlier, although significant flaws remain.

Understanding of the ISOs, on the other hand, remains disappointingly limited. Many simple models of the most widely studied ISO mode, the eastward-propagating Madden Julian oscillation¹¹ (MJO), have been constructed in attempts to reveal its fundamental mechanisms^{12–14}, but there are fundamental differences between them and little agreement on which of them, if any, is correct. The MJO is arguably the most significant mode of atmospheric variability at any subdecadal timescale that remains so unsatisfactorily explained. Why are there coherent modes of

variability on the intraseasonal timescale? What is the energy source? Why are they planetary in scale and why does the MJO propagate eastward at around 5 m s^{-1} ?

We focus here on the energy source, arguing that evidence, some but not all of it recent, suggests that feedbacks involving turbulent and radiative heat transfer between ocean and atmosphere drive the ISOs.

SURFACE HEAT FLUXES AS THE ENERGY SOURCE OF THE ISOs

The idea that air–sea interaction might provide the energy source for the MJO was first proposed over two decades ago^{15,16}. These authors used the quasi-equilibrium framework, which states that cumulus convection, owing its existence to the buoyancy of warm moist air rising from near the surface, acts quickly to eliminate that buoyancy, either by cooling and drying the surface or warming the upper atmosphere, or both^{17–19}. Quasi-equilibrium predicts that large-scale disturbances will not develop spontaneously owing to convection alone. However, if a basic-state easterly flow is assumed, quasi-equilibrium predicts that eastward propagating moist Kelvin wave disturbances—modified versions of the dry equatorial Kelvin waves which are fundamental linear modes of a shallow fluid layer on a rotating sphere—will develop owing to another process, namely variations in surface heat fluxes induced by the surface wind variations associated with the disturbance itself. This mechanism was termed wind–evaporation feedback or wind-induced surface heat exchange (WISHE). Disturbances can also develop spontaneously in models of this type (and in more complex models) from the interaction of clouds with electromagnetic radiation²⁰. Because of competing effects on the short-wave (solar) and long-wave (terrestrial) parts of the radiation spectrum²¹, deep cumulus clouds result in little net loss or gain of energy from the atmosphere to space. Their primary impact is a transfer of energy from ocean to atmosphere, analogously to a surface turbulent flux²².




Figure 1 Intraseasonal rainfall variance in the 30–90 day frequency band from the TRMM 3B42 data set (1998–2006). **a**, November–April. **b**, May–October. Bandpass filtering to retain only intraseasonal frequencies was done on daily data using two 60-point non-recursive digital filters with half-power points at 30 and 90 days.

In short, the simplest quasi-equilibrium models predict the emergence of large-scale weather disturbances such as ISOs only when convection interacts with processes, such as turbulent surface fluxes or radiation, that are moist diabatic. A moist diabatic process is one that alters the total entropy (including the component associated with the presence of water substance) of air parcels. This prediction of quasi-equilibrium is in contrast to so-called conditional instability of the second kind (CISK) models²³, which predict the emergence of large-scale disturbances without surface fluxes or radiation. For the MJO, frictional CISK in particular^{24,25}, in which near-surface circulations driven by surface friction have a crucial role in organizing convection, is a prominent theory that does not involve diabatic processes.

Observations indicate the need for modification of the original WISHE theory in several respects. The MJO is most strongly manifest in regions of mean surface westerlies, and the active phases occur in regions where the total surface wind is westerly^{26,27}, whereas both mean winds and total surface winds in active phases are easterly in the theory. In addition, spectral analysis²⁸ shows that the MJO is a distinct phenomenon from the ‘convectively coupled Kelvin waves’ that the original WISHE theory predicted. Although these observations are at odds with the original WISHE models in detail, they do not disprove the general notion that moist diabatic processes, and specifically radiative and turbulent heat exchange between the ocean and atmosphere, are integral to the ISOs. On the latter issue, the current state of understanding is unclear. We argue here that a substantial and growing body of evidence suggests a fundamental role for air–sea heat transfer in driving the ISOs.

LAND–SEA CONTRASTS IN INTRASEASONAL VARIABILITY

Over the oceans, the ISOs are associated with large variations in net heat flux. The flux from ocean to atmosphere during an active phase is greater than that during a suppressed phase by as much as 100 W m^{-2} or more, of which comparable fractions

are radiative and turbulent²⁹. These flux anomalies are temporally well correlated with precipitation anomalies in those oceanic regions where ISO convective variance is large³⁰, and at any given time within those regions the total surface flux is spatially correlated with precipitation. As disturbances propagate eastward or northward, the peak surface flux tends to lag the peak precipitation slightly, indicating that the fluxes are not responsible for the disturbances’ propagation even though they may well be essential to their existence.

Over land, the total net heat flux, and thus also its intraseasonal anomalies, are constrained to be small owing to the small effective heat capacity of the land surface. If net surface heat flux variations were important in driving intraseasonal variability in deep convection, the latter variability would be expected to be smaller over land than ocean. This is indeed the case, as shown in Fig. 1, produced using the Tropical Rainfall Measuring Mission (TRMM) 3B42 (ref. 31) data set from 1998 to 2006. Qualitatively similar results are found from other data sets with longer records (not shown), both for precipitation and related quantities such as top-of-atmosphere outgoing long-wave radiation, which is related to high cloudiness^{32–34}.

Climatological mean precipitation, although having similar large-scale structure to that of the variance, does not show the same land–sea contrasts on small scales (Fig. 2). This is so particularly in Southern Hemisphere summer, when climatological rainfall tends to maximize over the large Indonesian islands³⁵. That the magnitude of the intraseasonal variability is not a simple function of the mean suggests that different mechanisms control the two. Orography presumably does influence both, as demonstrated by the tendency of both mean and variance to maximize just to the west (and thus upstream, in westerly monsoon flow) of coastal mountain ranges, especially in northern summer³⁶. Nonetheless, we argue that differential surface heat capacity provides a more general explanation than orographic effects for the land–sea contrast in intraseasonal precipitation variance.




Figure 2 Climatological mean rainfall from the TRMM 3B42 data set (1998–2006). **a**, November–April. **b**, May–October.

Given the large spatial scales of the ISOs as usually defined, the relatively small-scale structures in the variance maps require some interpretation. Other observations have been interpreted to suggest that intraseasonal variations in convection at a point can be thought of as resulting from a local recharge–discharge oscillation^{33,37,38}; energy accumulates in the ocean mixed layer and lower troposphere while convection is suppressed, and is then vented to higher altitudes and surrounding regions when convection is active. A simple single-column model based on an implementation of quasi-equilibrium (ref. 39) and parameterized large-scale atmospheric dynamics captures this behaviour²². This model generates spontaneous intraseasonal oscillations if air–sea flux feedbacks are sufficiently strong. For weaker feedbacks, oscillations do not occur spontaneously, but forcing on intraseasonal timescales is effective in generating a response. We can think of the large-scale, propagating ISO as providing external forcing, and the single-column dynamics, dependent on the surface properties, as determining the amplitude of the local convective response. Whether free or forced, insufficient surface heat capacity (less than that of a few metres of water) prevents the model from sustaining significant intraseasonal variability, implying a prediction of small intraseasonal variance in convection over land, as observed.

Although convection at a point responds to forcing from the large-scale ISO, the aggregate response of convection over large horizontal areas must sustain the ISO, as the associated convective heating drives the large-scale flow perturbations. Thus, the mechanisms responsible for small-scale variance structure are presumably also relevant to the dynamics of the large-scale ISO itself.

USING MODELS TO TEST MECHANISMS

In numerical models, an individual physical process can be modified or eliminated, enabling direct assessment of the role of that process in the model. Turbulent and radiative fluxes are

computed by parameterizations contained in distinct portions of model code. By replacing the turbulent or radiative fluxes obtained from these parameterizations by specified values that do not depend on the model state (ideally, defined instead by annually and spatially varying climatologies computed from long control model integrations), the dynamical feedbacks involving those fluxes are disabled. The same feedbacks can also be disabled by eliminating the heat capacity of the ocean surface, rendering it a ‘swamp’. Comparing the ISO in a model with surface flux feedbacks removed with that in the control version of the same model gives a direct estimate of the role of those feedbacks in generating the control model’s ISO (ref. 16).

Calculations testing the role of surface turbulent flux feedbacks and radiative feedbacks in this manner have been carried out with a number of models, yielding mixed results. As model simulations of ISOs improve, however, such calculations become more compelling as guides to our understanding of the observed ISOs. In one recent study using a model in which the MJO simulation was state-of-the-art (although still flawed in some respects), the simulated MJO was strongly weakened when surface flux feedbacks were disabled⁴⁰. Similar results (not shown) have been obtained with a recent version of the National Oceanic and Atmospheric Administration Geophysical Fluid Dynamics Laboratory climate model. WISHE has also been shown to be important in an intermediate complexity model, where it acted synergistically with excitation by extratropical storms to generate an ISO (ref. 41).

Although suggestive, results from a small number of models are not entirely convincing. A more comprehensive set of calculations of this sort with a much larger number of models might enable a real step forward in our understanding of ISO dynamics, particularly if similar results are obtained from a large subset of the models, or from the subset in which ISOs most resemble the observed ones. Tests of this kind would be particularly interesting if conducted with recently developed global cloud-resolving models⁴² and the multiscale modelling framework^{43,44}, both of which show

exciting promise in the simulation of the ISOs. Surface turbulent momentum fluxes can also be evaluated in the same way as heat fluxes, enabling a direct test of the frictional CISK hypothesis^{24,25}. It is of course possible that the ISOs result from a combination of surface fluxes and moist adiabatic mechanisms such as frictional CISK, rather than one by itself.

In the case of the northward-propagating northern summer ISO (ref. 45), general circulation models have been used even less (compared with the MJO) to assess the role of distinct physical processes. In at least one model of intermediate complexity that gives a robust simulation of the northward-propagating ISO, surface flux feedbacks, specifically WISHE, are demonstrably important to the linear instability that produces the mode^{46,47}. The northward propagation seems to be controlled by large-scale dynamics^{46–48}.

OCEAN COUPLING

In addition to influencing the atmosphere, intraseasonal surface heat flux perturbations also induce sea surface temperature anomalies. To the extent that these anomalies then influence the atmosphere, the ISOs are coupled ocean–atmosphere modes. A number of studies have investigated this possibility, as summarized in recent reviews^{12–14}. Most (but not all) of these studies are consistent with the view that ocean coupling provides a modest amplification to the ISOs, and slows their propagation^{49,50}. The coupling question merits further study, but is secondary to that of the role of surface fluxes. If ocean coupling is important, surface fluxes must also be, because it is only through surface fluxes that the ocean can influence the atmosphere. The converse is not true; surface fluxes can be important while ocean coupling is not, because surface fluxes can vary strongly even if sea surface temperature remains constant. Thus, the role of fluxes is the more fundamental issue.

A PATH TOWARDS SOLVING THE ISO PROBLEM

We have argued that available evidence suggests an important, perhaps fundamental role for atmosphere–ocean heat fluxes, both turbulent and radiative, in driving the ISOs. If this were to be proved correct, it would still not constitute a theory for the ISOs, as the oscillations’ scales and structure would still remain to be explained. We would, however, be able to rule out the subset of existing theories in which only moist adiabatic dynamics are involved. Conversely, if surface heat fluxes were to be proved unimportant to the ISOs, we could focus our attention on moist adiabatic theories. Either situation would represent an improvement over the current state of confusion. The division into diabatic and adiabatic mechanisms is justified by analogy to extratropical atmospheric dynamics, where dry adiabatic models of geophysical fluid dynamics have been very successful. The land–sea contrast in intraseasonal precipitation and outgoing long-wave radiation variance also suggests this division, because it would be consistent with an important role for atmosphere–ocean heat exchange.

Numerical models have now improved in their simulations of the ISOs to the point that they may, collectively, enable a more convincing test of specific mechanisms, such as those involving surface fluxes. Reaching a conclusive outcome, if possible at all, requires the execution of simulations designed specifically for this purpose with a large number of models. Carrying out the necessary simulations is straightforward, but can require significant effort, depending on the details of the model code and the user’s facility with it. We hope our arguments help to convince modellers that this effort is worthwhile.

doi:10.1038/ngeo312

Published online: 21 September 2008.

References

1. Bessafi, M. & Wheeler, M. C. Modulation of south Indian Ocean tropical cyclones by the Madden–Julian oscillation and convectively-coupled equatorial waves. *Mon. Weath. Rev.* **134**, 638–656 (2006).
2. Mo, K. The association between intraseasonal oscillations and tropical storms in the Atlantic basin. *Mon. Weath. Rev.* **128**, 4097–4107 (2000).
3. Jones, C., Waliser, D. E., Lau, K. M. & Stern, W. The Madden Julian oscillation and its impact on northern hemisphere weather predictability. *Mon. Weath. Rev.* **132**, 1462–1471 (2004).
4. Bond, N. A. & Vecchi, G. A. On the Madden Julian oscillation and precipitation in Oregon and Washington. *Weath. Forecast.* **18**, 600–613 (2003).
5. McPhaden, M. J. The genesis and evolution of the 1997–98 El Niño. *Science* **283**, 950–954 (1999).
6. Webster, P. J. & Hoyos, C. Prediction of monsoon rainfall and river discharge on 15–30 day time scales. *Bull. Am. Meteorol. Soc.* **85**, 1745–1765 (2004).
7. Waliser, D. et al. The experimental MJO prediction project. *Bull. Am. Meteorol. Soc.* **87**, 425–431 (2006).
8. Vitart, F., Woolnough, S., Balmased, M. A. & Tompkins, A. M. Monthly forecast of the Madden-Julian oscillation using a coupled GCM. *Mon. Weath. Rev.* **135**, 2700–2715 (2007).
9. Zhang, C. et al. Simulations of the Madden-Julian Oscillation in four pairs of coupled and uncoupled models. *Clim. Dyn.* **27**, 573–592 (2006).
10. Lin, J. et al. Tropical intraseasonal variability in 14 IPCC AR4 climate models. Part I: Convective signals. *J. Clim.* **19**, 2665–2690 (2006).
11. Madden, R. A. & Julian, P. R. Observations of the 40–50 day tropical oscillation—A review. *Mon. Weath. Rev.* **122**, 814–837 (1994).
12. Zhang, C. Madden–Julian oscillation. *Rev. Geophys.* **43**, doi:10.1029/2004RG000158 (2005).
13. Wang, B. in *Intraseasonal Variability in the Atmosphere–Ocean Climate System* (eds Lau, W. K. M. & Waliser, D. E.) 307–360 (Praxis Publishing, Berlin, 2005).
14. Waliser, D. E. in *The Asian Monsoon* (ed. Wang, B.) 203–257 (Springer-Praxis, Berlin, 2006).
15. Emanuel, K. A. An air–sea interaction model of intraseasonal oscillations in the tropics. *J. Atmos. Sci.* **44**, 2324–2340 (1987).
16. Neelin, J. D., Held, I. M. & Cook, K. H. Evaporation–wind feedback and low-frequency variability in the tropical atmosphere. *J. Atmos. Sci.* **44**, 2341–2348 (1987).
17. Arakawa, A. & Schubert, W. H. Interaction of a cumulus cloud ensemble with the large-scale environment, Part I. *J. Atmos. Sci.* **31**, 674–701 (1974).
18. Emanuel, K. A., Neelin, J. D. & Bretherton, C. S. On large-scale circulations in convecting atmospheres. *Q. J. R. Meteorol. Soc.* **120**, 1111–1143 (1994).
19. Arakawa, A. The cumulus parameterization problem: Past, present and future. *J. Clim.* **17**, 2493–2525 (2004).
20. Fuchs, Z. & Raymond, D. J. Large-scale modes of a nonrotating atmosphere with water vapor and cloud-radiation feedbacks. *J. Atmos. Sci.* **59**, 1669–1679 (2002).
21. Ramanathan, V. et al. Cloud-radiative forcing and climate: Results from the Earth radiation budget experiment. *Science* **243**, 57–63 (1989).
22. Sobel, A. H. & Gildor, H. A simple time-dependent model of SST hot spots. *J. Clim.* **16**, 3978–3992 (2003).
23. Lindzen, R. S. Wave–CISK in the tropics. *J. Atmos. Sci.* **31**, 156–179 (1974).
24. Wang, B. Dynamics of tropical low-frequency waves: An analysis of the moist Kelvin wave. *J. Atmos. Sci.* **45**, 2051–2065 (1988).
25. Moskowitz, B. M. & Bretherton, C. S. An analysis of frictional feedback on a moist equatorial kelvin mode. *J. Atmos. Sci.* **57**, 2188–2206 (2000).
26. Kiladis, G. N., Meehl, G. A. & Weickmann, K. M. Large-scale circulation associated with westerly wind bursts and deep convection over the western equatorial Pacific. *J. Geophys. Res.* **99**, 18527–18544 (1994).
27. Zhang, C. & McPhaden, M. Intraseasonal surface cooling in the equatorial western Pacific. *J. Clim.* **13**, 2261–2276 (2000).
28. Wheeler, M. & Kiladis, G. N. Convectively coupled equatorial waves: Analysis of clouds and temperature in the wavenumber–frequency domain. *J. Atmos. Sci.* **56**, 374–399 (1999).
29. Shinoda, T., Hendon, H. H. & Glick, J. Intraseasonal variability of surface fluxes and sea surface temperature in the tropical western Pacific and Indian oceans. *J. Clim.* **11**, 1685–1702 (1998).
30. Araligidad, N. M. & Maloney, E. D. Wind-driven latent heat flux and the intraseasonal oscillation. *Geophys. Res. Lett.* **35**, L04815 (2008).
31. Adler, R. F., Huffman, G. J., Bolvin, D. T., Curtis, S. & Nelkin, E. J. Tropical rainfall distributions determined using TRMM combined with other satellite and rain gauge information. *J. Appl. Meteorol.* **39**, 2007–2023 (2000).
32. Zhang, C. D. & Hendon, H. H. Propagating and standing components of the intraseasonal oscillation in tropical convection. *J. Atmos. Sci.* **54**, 741–752 (1997).
33. Fasullo, J. & Webster, P. J. Warm pool SST variability in relation to the surface energy balance. *J. Clim.* **12**, 1292–1305 (1999).
34. Sperber, K. R. Madden–Julian variability in NCAR CAM2.0 and CCSM2.0. *Clim. Dyn.* **23**, 259–278 (2004).
35. Qian, J.-H. Why precipitation is mostly concentrated over islands in the maritime continent. *J. Atmos. Sci.* **62**, 1428–1441 (2008).
36. Hoyos, C. D. & Webster, P. J. The role of intraseasonal variability in the nature of Asian monsoon precipitation. *J. Clim.* **20**, 4402–4424 (2007).
37. Waliser, D. E. Formation and limiting mechanisms for very high sea surface temperature: Linking the dynamics and the thermodynamics. *J. Clim.* **9**, 161–188 (1996).
38. Stephens, G. L., Webster, P. J., Johnson, R. H., Engelen, R. & L’Ecuyer, T. Observational evidence for the mutual regulation of the tropical hydrological cycle and tropical sea surface temperatures. *J. Clim.* **17**, 2213–2224 (2004).
39. Neelin, J. D. & Zeng, N. A quasi-equilibrium tropical circulation model—formulation. *J. Atmos. Sci.* **57**, 1741–1766 (2000).
40. Maloney, E. D. & Sobel, A. H. Surface fluxes and ocean coupling in the tropical intraseasonal oscillation. *J. Clim.* **17**, 4368–4386 (2004).
41. Lin, J. W.-B., Neelin, J. D. & Zeng, N. Maintenance of tropical intraseasonal variability: Impact of evaporation–wind feedback and midlatitude storms. *J. Atmos. Sci.* **57**, 2793–2823 (2000).
42. Miura, H., Satoh, M., Nasuno, T., Noda, A. T. & Oouchi, K. A Madden–Julian oscillation event realistically simulated by a global cloud-resolving model. *Science* **318**, 1763–1765 (2007).
43. Grabowski, W. W. MJO-like coherent structures: Sensitivity simulations using the cloud-resolving convection parameterization (CRCP). *J. Atmos. Sci.* **60**, 847–864 (2003).
44. Khairoutdinov, M., DeMott, C. & Randall, D. A. Evaluation of the simulated interannual and subseasonal variability in an AMIP-style simulation using the CSU multiscale modeling framework. *J. Clim.* **21**, 413–431 (2008).

45. Goswami, B. N. in *Intraseasonal Variability in the Atmosphere–Ocean Climate System* (eds Lau, W. K. M. & Waliser, D. E.) 19–61 (Springer, Berlin, 2005).
46. Bellon, G. & Sobel, A. H. Poleward-propagating intraseasonal monsoon disturbances in an intermediate-complexity axisymmetric model. *J. Atmos. Sci.* **65**, 470–489 (2008).
47. Bellon, G. & Sobel, A. H. Instability of the axisymmetric monsoon flow and intraseasonal oscillation. *J. Geophys. Res.* **113**, D07109 (2008).
48. Drbohlav, H.-K. & Wang, B. Mechanism of the northward-propagating intraseasonal oscillation: Insights from a zonally symmetric model. *J. Clim.* **18**, 952–972 (2005).
49. Klingaman, N. P., Inness, P. M., Weller, H. & Slingo, J. M. The importance of high-frequency sea-surface temperature variability to the intraseasonal oscillation of Indian Monsoon Rainfall. *J. Clim.* **21**, doi:10.1175/2008JCLI2329.1 (2008).
50. Bellon, G., Sobel, A. H. & Vialard, J. Ocean-atmosphere coupling in the monsoon intraseasonal oscillation: A simple model study. *J. Clim.* **21**, doi:10.1175/2008JCLI2305.1 (2008).

Acknowledgements

This paper was written while the first author was on sabbatical at the Australian Bureau of Meteorology, and he thanks M. Wheeler, H. Hendon, H. Zhu, S. Cleland and L. Chappell for stimulating discussions on the MJO. This work was supported by the Climate and Large-Scale Dynamics Program of the National Science Foundation under grants ATM-0832868 (E.D.M.) and ATM-054273 (A.H.S.), by the Precipitation Measurement Mission program of the National Aeronautics and Space Administration under grant NNX07AD21G (A.H.S.) and by award NA05OAR4310006 from the National Oceanic and Atmospheric Administration, US Department of Commerce (E.D.M.). The statements, findings, conclusions and recommendations do not necessarily reflect the views of NSF, NASA, NOAA or of the Department of Commerce.

Author information

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions>. Correspondence and requests for materials should be addressed to A.H.S.